
STRESS COPING AND INFECTIUS ILLNESS: PERSISTENTLY LOW NATURAL KILLER CELL (U) PITTSBURGH UNIV PA S M LEVY ET AL 14 MAR 88 N00014-87-K-0224 AD-8192 826 F/G 5/8 NL UNCLASSIFIED



(U) SECURITY CLASSIFICATION OF THIS PAGE	TIC	ហ	16 FILE	Copy	
SECURITY CLASSIFICATION OF THIS PAGE	E COMEN A	N PAGE		1 1	Form Approved OMB No. 0704-0188
4	? 2 5 1988	1b. RESTRICTIVE	MARKINGS		
-		NA NA	/AVAILABILITY OF	PLOOP	
AD-A192 826	Z-D		ion Unlimit		i
- FERFORMUNG ORGANIZATION REPORT NUMBER		S MONITORING	ORGANIZATION RE	OORT NUMB	ER/C)
University of Pittsburgh	EN(3)	NA NA	SKIMIZATION RE	FORT NOINE	EN(3)
6a NAME OF PERFORMING ORGANIZATION	66 OFFICE SYMBOL		ONITORING ORGAN	NIZATION	
University of Pittsburgh	(If applicable) ONR		f Naval Res		
6c. ADDRESS (City, State, and ZIP Code)		7b. ADDRESS (Cit	y, State, and ZIP C	ode)	
200 Meyran Ave., 2nd Floor			uincy Stree		
Pittsburgh, PA 15213		Arlington	n, VA 2221	7-5000	
8a NAME OF FUNDING/SPONSORING	8b. OFFICE SYMBOL	9. PROCUREMENT	T INSTRUMENT IDE	NTIFICATION	NUMBER
ORGANIZATION Office of Naval Research	(If applicable) ONR	N00014-8	7-K-0224		i
8c. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street	<u> </u>	10. SOURCE OF F	UNDING NUMBERS	S	
Arlington, A 22217-5000		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.
A 11119 (611, 11 2221) 3000		61153N	RR04108	R&T442d	
11 TITLE (Include Security Classification)		<u></u>	L	L	
''Stress, Coping, and Infecti	us Illness: P	ersistently	Low Natural	Killer	Cell Activity
as a Host Risk Factor"					
12 PERSONAL AUTHOR(S) Sandra M. Levy, Ph.D., Ronald	R Herberman	M.D. Theres	a Whiteside	Ph.D.	Anne Simons.
13a TYPE OF REPORT		14. DATE OF REPO			
	/87 TO 2/29/88				
16 SUPPLEMENTARY NOTATION					
17 COSATI CODES	18. SUBJECT TERMS (Continue on reverse	e if necessary and	identify by	block number)
FIELD GROUP SUB-GROUP	Key words:		-		ity, fatigue,
08] ((0) ((0) (0) (0)				tal distress.
	<u> </u>				
Recent Japanese research, as					
have pointed to a possible a	ssociation bet	ween mood and	d lowered na	atural k	iller (NK)
cell activity. In this prev persistently low NK activity					
more serious illness on foll					
approximately, sixty-two nor	mal individual	s to this pro	ospective p	roiect.	Subjects are
assessed both psychologicall					
over a six-month follow-up p	eriod in order	to identify	predictors	of infe	ctious illness
over time. As in the prelim	inary work, we	are finding	that approx	ximately	1/3 of the
sample has persistently low	NK activity, a	nd this immu	nological pa	attern i	s strikingly
associated with daily stress	levels, as we	II as reports	s of depress	sion and	chronic
anxiety. In a very prelimin between this low NK activity	ary rashion, w	e nave also	iound a trem	nd OT as	SUCTATION Iness
	pattern, and				111633.
O DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED SAME AS R		21. ABSTRACT SEC	CURITY CLASSIFICA	TION	
Dr. I. A. Hajde (or other ON		226, TELEPHONE (1 (202) 696	nclude Area Code) - 4055	22c. OFFIC	RSYMBOL

DD Form 1473, JUN 86

Dr. I. A. 'lajde (or other ONR Sci. Off.)

CONTRACTOR CONTRACTOR

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

ANNUAL REPORT

Stress, Coping, and Infectious Illness: Persistently Low Natural Killer Cell Activity as a Host Risk Factor

ONR Contract #: N00014-87-K-0224

I. Introduction

The study of stress and coping responses linked with disease end-points in animal systems has revealed enhancement of disease risk under a variety of stress conditions (Riley, 1981; Visintainer, Volpicelli, and Seligman, 1982; Weiss, 1972; Laudenslager, Ryan, Drugan, et al., 1983). A number of such studies have also examined endocrinological and immunological mechanisms potentially mediating behavior and disease end-points (Shavit, Lewis, Terman, et al., 1983; Sklar and Anisman, 1979; Greenberg, Dyck, and Sandler, 1985; Schneiderman, 1986). However, there are scant clinical data demonstrating a link between distress response and disease end-points mediated by regulatory mechanisms such as immune function.

A number of studies have demonstrated a correlation between stressful life events of various kinds and increased incidence of acute, infectious illnesses, such as upper respiratory infections and infectious mononucleosis, as well as outbreaks of herpes simplex (Ishigami, 1919; Hinkle and Plummer, 1952; McClelland, Alexander, and Marks, 1982; Kemeny, et al., 1986). Several investigations have also indicated an association between stressful events and lymphocyte alteration (Palmblad, Blomback, Egberg, et al., 1977; Bartrup, Luckhurst, Lazarus, et al., 1977; Crary, Borysenko, Sutherland, et al., 1983; Jemmott and Locke, 1984). However, with only a few exceptions (Kasl, et al., 1979 and Kemeny, et al., 1986), a major limitation of most human studies has been that they were not conducted in a prospective fashion, examining the association of life stress and coping ability with hormonal and immunological changes, with subjects then followed to assess the incidence of disease episodes. Since most studies have not made the final, longitudinal link with actual disease, the biologic significance of stress-related immune impairment remains unclear.

Preliminary work by our research team suggests that it may be possible to identify a subgroup of vulnerable individuals at particular risk of infectious illness. For example, research carried out by Aoki, Herberman and colleagues (1985) points to a possible association between mood and lowered NK activity. They identified a subgroup within a patient sample characterized by low NK activity, remittent fever, and self-reported depression and fatigue. The depressive and fatigue-like symptoms were sufficiently prominant that these individuals had frequently been seen by psychiatrists rather than by other health care specialists. These investigators concluded that these individuals may be suffering from a new immunological disorder termed Low Natural Killer Cell Syndrome (LNKS), and reported some success in treating them with an immunopotentiator, letinen.

The Principal Investigator and Co-Principal Investigator on this proposed project (S. Levy and R. Herberman) conducted preliminary work in the

Biological Response Modifiers Program at the NCI with resultant data consistent with Aoki, et al.'s. That study examined the predictive value of daily stressors, personality, and coping factors, as well as repeated baseline measures of natural immunity (NK activity) and hormonal distress markers (excreted epinephrine and norepinephrine) relevant to episodes of infectious illness in a sample of healthy laboratory volunteers. Results from that prospective study revealed a subgroup at risk for disease, identified as having "low natural killer cell syndrome" (LNKS). Such individuals, operationally defined as having persistently low functional levels of NK activity across all three times of measurement (three baseline measures separated by two-week intervals), tended to report more serious illness over a three-month follow-up When comparing the psychological and demographic profile of the LNKS versus "normal" NK group, the LNKS individuals tended to be younger, report more daily distress, and had higher levels of excreted urinary catecholamines than individuals with normal NK activity.

Taken together, these studies suggest that inadequate coping (reflected in reports of depression and/or fatigue, coupled with report of high levels of daily hassles), psychological distress (reflected in higher levels of norepinephrine and epineprine excretion) and immune function (specifically NK activity) may interact to increase risk for infectious illness.

The direction of association is unclear at this time. It is plausible that psychological distress produced by inadequate problem solving of life's hassles may compromise immune function and lead to the development of illness. On the other hand, low levels of natural immunity may produce cognitive and behavioral effects such as dysphoria and inability to cope. This project is designed to assess the relevant variables over a sufficient length of time to begin to understand these complex relationships.


II. Progress Report

CONTRACTOR OF THE PROPERTY OF

During the first eleven months of this contract, we have trained a research associate who has served as a key project staff member for recruiting individuals from the community. We have also trained two lab technicians, and have begun carrying out endogenous opioid assays (plasma beta endorphin) in Dr. Fernstrom's laboratory, and have begun analyzing NK activity, and conducting flow cytometry analyses in Dr. Whiteside's laboratory.

We have accrued approximately sixty normal community volunteers to date Initially, we intended to accrue both normal for this prospective study. volunteers, and individuals who received a psychiatric diagnosis of dysthymia (chronic low level depression). However, it quickly became clear that the latter category of subjects was not available for study, as most wished to receive some sort of psychological treatment. We initially thought that we would randomize half of this sub-sample to a treatment and a wait-list control However, after much consideration, we decided that the latter option would take us too far afield from the intention of this current project. Hence, in consultation with our project officer, we decided to focus on accruing a normal sample, from which the findings would be most generalizable to a military population. We have had very good response from the community, and our accrual of this sample to date has been excellent.





q

<u>Demographics</u>. The mean age of the sample accrued so far is 30 (range, 18-44 years). The average educational level is 17.5 years, with approximately 60% of the sample as female. Fifty-three percent of the sample is single, with 33% married, and 13% divorced.

In terms of NK function, mean NK activity at the 50:1 E/T ratio = 44% (range, 18-80%), and the mean lytic unit value is 131.6. Approximately 24% of the sample shows persistently low NK activity, defined as NK function below the group mean at baseline, and also at three follow-up visits.

Follow-up illnesses reported. The following categories of illnesses, with numbers of individuals reporting having experienced these health end-points, are shown in Table 1.

TABLE I Illness reported on Follow-up

	inness reported on ronow-up
Illness Category	Numbers of Individuals
Cold	22
Influenza	4
Pneumonia	0
Cold sore	4
Gum infection	6
Mononucleosis	0
Strep throat	3
Gastrointestinal virus	9
Fever symptoms	7
Sore throat	7
Other	25

Preliminary Pearson correlations between NK activity and other study key Table 2 displays selected correlations between study variables of variables. major interest in this project. As can be seen, there are some interesting clusters of correlations in this Table. First, NK activity, at two E/T ratios, as well as measured in lytic units, was significantly negatively correlated with reported days ill with strep infection. (Please note that the N reporting strep infection is small, and this correlation should be viewed with caution.) function was also negatively correlated with frequency and severity of daily Hassles, or daily stressors, are also significantly stressors or hassles. positively correlated with other indices of distress, namely total Profile of Mood States scores, and specifically, with the report of fatigue-like symptoms. The latter is particularly interesting because of Aoki, et al's., (1985) findings that fatigue, or "uncomfortable general duliness" is associated with what he terms Low NK Syndrome. (Please note: We have not analyzed the largest illness category of "Other," so these data are quite preliminary in terms of simple associations with illness episodes.) Interestingly, NK function was positively correlated with age, and this is consistent with findings from other studies that we have carried out.

Persistently low NK activity. As indicated earlier, approximately 29% of the subjects had persistantly low NK activity at baseline and at two follow-up periods, and 24% of subjects has persistently low NK activity at three repeat

assessment periods (baseline, and three follow-up measurements). latter as the most stringent criterion for persistently low NK activity, we carried out a Chi-Square test, stratifying subjects into low NK activity versus other, and serious versus non-serious illness on follow-up (omitting the "Other" category, and the symptom category of fever). The "serious" category included: Influenza, pneumonia, mononucleosis, strep infection. The "non-serious" illness category included: gastrointestinal virus. Cold. cold sore, gum infection, and minor sore throat. A non-significant trend was found for this preliminary analysis, with those falling into the persistently low NK category tending to report more serious illness on follow-up (Likelihood ratio Chi-Square = 1.5, p<.2).

When comparing those whith persistently low NK activity with others in the sample, many of our earlier pilot findings were confirmed here. Individuals with the low NK profile were also significantly more chronically anxious (t=2.1, p<.05), more depressed (t=1.7, p<.04), more confused (t=1.9, p<.05), with trends in the direction of more fatigue (t=1.5, p<.1) and acute anxiety (t=1.6, p<.1). Again, those with persistently low NK activity reported strikingly more frequent daily stressors (t=2.6, p<.02), more severe stressors (t=2.8, p<.01) and more intense daily stress levels (t=2.6, p<.01).

In sum, we are finding very similar trends and patterns in the current study to those we found in our pilot work for this project. It should be emphasized that these are preliminary data, and that when this prospective study is complete, we will be able to test by causal modeling procedures the direction and strength of associations between our predictor variables and illness endpoints.

REFERENCES

Aoki, T., Usuda, Y., Miyakoski, H., Tamura, K., Fujimoto, M., and Herberman, R., (1985). Low NK syndrome (LNKS). Proceedings of the Seventh Symposium on Host Defense Mechanisms Against Cancer, Hakone, November 8-10, 1985.

Bartrup, R., Lazarus, L., Luckhurst, E., Kiloh, L., and Penny, R., (1977). Depressed lymphocyte function after bereavement. <u>Lancet</u>, <u>1</u>, 834-836.

Crary, B., Borysenko, M., Sutherland, D., Kutz, I., Borysenko, J., and Benson, H., (1983). Decrease in mitogen responsiveness of mononucelar cells from peripheral blood after epinephrine administration in humans. <u>Journal of Immunology</u>, 130, 604-607.

Greenberg, A., Dyck, D., and Sandler, L., (1984). Opponent processes, neurohormones, and natural resistance. In B. Fox and B. Newberry (Eds.), <u>Psychoneuroendocrine Systems in Cancer and Immunity</u>. Toronto: Hogrefe.

Hinkle, L., and Plummer, N., (1952). Life stress and industrial absenteeism. Industrial Medicine and Surgery, 21, 363-375.

Ishigami, T., (1919). The influence of psychic acts on the progress of pulmonary tuberculosis. <u>American Review of Tuberculosis</u>, 2, 470-484.

Jemmott, J., and Locke, S., (1984). Psychosocial factors, immunologic mediation, and human susceptibility to infectious diseases: How much do we know? Psychological Bulletin, 95, 78-108.

Kasl, S., Evans, A., and Neiderman, J., (1979). Psychosocial risk factors in the development of infectious mononucleosis. <u>Psychosomatic Medicine</u>, 41, 445-466.

Kemeny, M., Cohen, F., and Zegams, L., (1986). The relationship of coping strategies to immunity and genital herpes recurrence. Paper presented to the Society of Behavioral Medicine Meeting, San Francisco, March 3-5.

Levy, S., (1987). Behavior, immunity, and breast cancer: The role of individual differences. Presented at the 95th Annual Convention of the American Psychological Association at New York City, August 28, 1987.

McClelland, D., Alexander, C., and Marks, E., (1982). The need for power, stress, immune function, and illness among male prisoners. <u>Journal of Abnormal Psychology</u>, 91, 61-70.

Palmblad, J., Blomback, M., Egberg, N., Froberg, J., Karlsson, C., and Levi, L., (1977). Experimentally induced stress in man: Effects on blood coagulation and fibrinolysis. <u>Journal of Psychosomatic Research</u>, 21, 87-92.

Schneiderman, N., (1986). Issues on intervention in coronary prone behavior. Paper presented at the Banff International Conference on Behavioral Science, Banff, Alberta, Canada, March 16-20.

Shavit, Y., Lewis, J., Terman, G., Gale, R., and Liebeskind, J., (1984). Opioid peptides mediate the suppressive effect of stress on natural killer cell cytotoxicity. <u>Science</u>, 223, 188-190.

Sklar, L., and Anisman, H., (1979). Stress and coping factors influence tumor growth. Science, 205, 513-515.

TABLE 2 Pearson Correlations Between Study Variables

					200		במו זמו כמו בומרומוז הבשבבו סרמה ימו ומחובז	מרחת אחו	90103					
	NK 50:1 25:1 LU AGE	25:1	3	AGE	COLD	FLU	STREP	FEVER	FAT.	TOT POMS Leu 19	Leu 19	H-FREQ.	H-SEV.	긒
NK (50:1)		***6.	**26*	.92** .46**	NS	NS	33*	NS	NS	NS	NS	36**	31**	NS
NK (25:1)			**96*	**55. **96.	NS	NS	25	NS	18	NS	NS	-,41**	40**	NS
NK Lu				.45**	NS	NS	25	NS	NS	NS	NS	-,32**	31**	NS
Age					51**	NS	NS	NS	NS		.20	NS	NS	NS
Cold						NS	NS	NS	NS		40**	NS	NS	NS
Flu							NS	NS	NS	NS	SN	NS	NS	NS
Strep								.31*	NS	NS	. SN	NS	NS	.28
Fever									NS	NS	NS	NS	NS	NS
Fatigue										.67**	NS	.28*	.34*	.44*
Tot POMS											NS	.48**	**95.	.66**
Leu 19												NS	NS	NS
Hassles-Freq.													.97	.27
Hassles-Sev.														. 44
														•

. p .05

DISTRIBUTION LIST

Behavioral Immunology Program

Annual, Final and Technical Reports (one copy each except as noted)

INVESTIGATORS

Dr. Itamar B. Abrass Department of Medicine University of Washington Harborview Medical Center Seattle, WA 98104

Dr. Prince K. Arora NICHD, Bldg 6, Room 132 National Institutes of Health Bethesda, MD 20892

Dr. Karen Bulloch Helicon Foundation 4622 Sante Fe Street San Diego, CA 92109

Dr. Michael D. Cahalan Department of Physiology and Biophysics University of California, Irving Irvine, CA 92717

Dr. Donald A. Chambers Health Sciences Center University of Illinois at Chicago P.O. Box 6998 Chicago, IL 60680

Dr. Christopher L. Coe Department of Psychology Harlow Primate Laboratory University of Wisconsin Madison, WI 53715

Dr. Walla L. Dempsey
Department of Microbiology and Immunology
The Medical College of Pennsylvania
3300 Henry Avenue
Philadelphia, PA 19129

Dr. Adrian J. Dunn
Department of Neurescience
University of Florida
College of Medicine
Gainsville, FL 32610

Dr. David L. Felten
Department of Anatomy
University of Rochester
School of Medicine
601 Elmwood Avenue
Rochester, NY 14642

Dr. John F. Hansbrough Department of Surgery UCSD Medical Center 225 Dickinson Street San Diego, CA 92103

Dr. William F. Hickey Neuropathology Laboratories 454 Johnson Pavilion University of Pennsylvania Philadelphia, PA 19104

Dr. Robert L. Hunter Department of Pathology Emory Univ. School of Medicine WMB 760 Atlanta, GA 30322

Dr. Terry C. Johnson Division of Biology Ackert Hall Kansas State University Manhattan, KS 66506

Dr. Sandra Levy University of Pittsburgh School of Medicine 3811 O'Hara Street Pittsburgh, PA 15213 Dr. Lester Luborsky
Department of Psychiatry
308 Piersol Building/G1
University of Pennsylvania Hospital
Philadelphia, PA 19104

Dr. Steven F. Maier Department of Psychology University of Colorado Campus Box 345 Boulder, CO 80309

Dr. Michael H. Melner Department of Biochemistry Univ of Miami School of Medicine 1600 N.W. 10th Avenue Miami, FL 33136

Dr. Vera B. Morhenn Department of Dermatology Stanford University Medical School Stanford, CA 94305

Dr. Jose R. Perez-Polo Gail Borden Bldg., Rm., 436 University of Texas Medical Branch Galveston, TX 77550-2777

Dr. Howard R. Petty Department of Biological Sciences Wayne State University Detroit, MI 48202

Dr. Bruce S. Rabin Clinical Immunopathology Childrens Hospital University of Pittsburgh Sch of Medicine Pittsburgh, PA 15213

Dr. Seymour Reichlin Director, Clinical Study Unit New England Midical Center Hospitals, Inc. 171 Harrison Avenue Boston, MA 02111 Dr. Eric M. Smith
Department of Psychiatry
University of Texas Medical Branch
Galveston, TX 77550

Dr. Arthur A. Stone Department of Psychiatry
State University of New York at Stony Brook
Stony Brook, NY 11794

Annual, Final and Technical Reports (one copy each except as noted)

ADMINISTRATORS

Dr. Jeannine A. Majde, Code 1141CB (2 copies) Scientific Officer, Immunology Program Office of Naval Research 800 N. Quincy Street Arlington, VA 22217-5000

Administrator (2 copies) (Enclose DTIC Form 50) Defense Technical Information Center Building 5, Cameron Station Alexandria, VA 22314

Administrative Contracting Officer
ONR Resident Representative
(address varies - obtain from business office)

Program Manager Biological/Human Factors Division Office of Naval Research, Code 125 800 N. Quincy Street Arlington, VA 22217-5000

Program Manager Support Technology Directorate Office of Naval Technology, Code 223 800 N. Quincy Street Arlington, VA 22217-5000

Annual and Final Reports Only (one copy each)

DOD ACTIVITIES

Commanding Officer Naval Medical Command Washington, DC 20372

Commanding Officer Naval Medical Research & Development Command National Naval Medical Center Bethesda, MD 20814

Director, Infectious Diseases Program Center Naval Medical Research Institute National Naval Medical Center Bethesda, MD 20814

Commander
Chemical and Biological Sciences Division
Army Research Office, P.O. Box 12211
Research Triangle Park, NC 27709

Commander
U.S. Army Research and Development Command
Attn: SGRD-PLA
Fort Detrick
Frederick, MD 21701

Final and Technical Reports Only

Director, Naval Research Laboratory (6 copies) Attn: Technical Information Division, Code 2627 Washington, DC 20375 Commander
USAMRIID
Fort Detrick
Frederick, MD 21701

Directorate of Life Sciences Air Force Office of Scientific Research Bolling Air Force Base Washington, DC 20332

Library
Armed Forces Radiation Research
Institute
Bethesda, MD 20814-5145

TE DA